



Patient Information and Consent To Treatment With Buprenorphine/Naloxone

Suboxone® (a tablet or strip with buprenorphine and naloxone in it) is an FDA approved medication for treatment of people with heroin or other opioid (prescription pain medication) addiction. Buprenorphine can be used for detoxification or for maintenance therapy. Maintenance therapy can continue as long as medically necessary. There are other treatments for opiate addiction, including methadone, naltrexone, and some treatments without medications that include counseling, groups and meetings.

If you are dependent on opiates - any opiates - **you should be in as much withdrawal as possible when you take the first dose of buprenorphine. It you are not in withdrawal, buprenorphine can cause severe opiate withdrawal.** For that reason, you should take the first dose in the office and remain in the office for at least 2-3 hours. You should not drive or operate machinery until you know how buprenorphine/naloxone affects you. Therefore, you must arrange not to drive after your first dose, because some patients get drowsy until the correct dose is determined for them. You must either take public transportation to the office visits for induction or you must have someone drive you to the induction appointment(s).

Some patients find that it takes several days to get used to the transition from the opiate they had been using to buprenorphine. During that time, any use of other opiates may cause an increase in symptoms. After you become stabilized on buprenorphine, it is expected that other opiates will have less effect. Attempts to override the buprenorphine by taking more opiates could result in an opiate overdose. You should not take any other medication without discussing it with the physician first.

Combining buprenorphine with alcohol or other sedating or tranquilizer medications is dangerous. The combination of buprenorphine with benzodiazepines (such as Valium®, Librium®, Ativan®, Xanax®, Klonopin®, etc.) has resulted in deaths. You must agree not to take such medications without discussing this with your doctor. If you think you need to take these medications you may be referred to other treatment for your opioid addiction and you agree to this before starting buprenorphine/naloxone.

Although buprenorphine given under the tongue has not been shown to be liver-damaging (except when people take very large amounts of buprenorphine or sometimes, if Hepatitis C is present), your doctor will monitor your liver tests while you are taking buprenorphine. (This is a blood test.)

The form of buprenorphine (Suboxone®) you will be taking is a combination of buprenorphine with a short-acting opiate blocker (naloxone) in a 4 to 1 ratio (4 mg of buprenorphine to 1 mg naloxone). **It will maintain physical dependence**, and if you discontinue it suddenly, you will likely experience withdrawal symptoms. If you are not already dependent, you should not take buprenorphine, it could eventually cause physical dependence.

Buprenorphine/naloxone tablets or the buprenorphine/naloxone strip must be held under the tongue until it is dissolved completely. You will be given your first dose at the clinic, and you will have to wait as it dissolves, and for two hours after it dissolves, to see how you react. **It is important not to talk and to swallow as little as possible until the medication dissolves.** This takes up to ten minutes.

Buprenorphine is then absorbed over the next 30 to 120 minutes from the tissue under the tongue. Buprenorphine is poorly absorbed from the stomach. **If you swallow the tablet, you will not have the important benefits of the medication, and it may not relieve your withdrawal.**

Most patients end up at a daily dose of 12/3-16/4 mg of buprenorphine. Beyond that dose, the effects of buprenorphine plateau, so there may not be any more benefit to increase in dose. It may take several weeks to determine just the right dose for you. The first dose is usually 2/0.5-4/1 mg.

If you are transferring to Suboxone® from methadone maintenance, your dose has to be tapered until you have been at or **below 30mg for at least a week**. There must be **at least 24 hours** (preferably longer) between the time you take your last methadone dose and the time you are given your first dose of buprenorphine. Your doctor will examine you for clear signs of withdrawal, and you will not be given buprenorphine until you are in withdrawal.

I have read and understand these details about buprenorphine treatment. I have had my questions about buprenorphine/naloxone and treatments for opioid addiction answered. I wish to be treated with buprenorphine.

Signature

Date

Witness

Date